

Comment II on J.M. Salazar and L. Brenig's contribution: “Computer Simulations and Kinetic Theory of an Inelastic Granular Gas”

M.H. Ernst

*Institute for Theoretical Physics, P.O.Box 80.006
3509 TA Utrecht, The Netherlands; E-mail:ernst@phys.uu.nl*

Abstract

In two recent articles Salazar and Brenig question the validity of kinetic theory for granular gases and fluids, on the basis of a supposedly exact hierarchy of coupled equations for the velocity moments, which the authors derive from the BBGKY-hierarchy. Their derivation contains several errors, which are exposed. Moreover, they support their findings with results from direct monte carlo simulation (DSMC) of the Boltzmann equation, which supposedly show that Haff's homogeneous cooling law only holds for times shorter than one mean free time. However, their DSMC results have no physical significance as they are carried out at a density, twice close packing.

The recent comment in cond-mat/9911276 by Y. Elskens is completely in line with my criticism, voiced at the workshop on Granular Gases, that took place in Bad Honnef on March 8 - 12, 1999 (see Ref. [1]). It prompted the publication of the notes, below made at that conference, which raise a number of additional objections.

Salazar and Brenig in Ref. [2] claim to have obtained the following analytic and simulation results for undriven (freely cooling) granular gases of Inelastic Hard Spheres (IHS):

- (i) A Direct Simulation Monte Carlo (DSMC) solution of the Boltzmann equation, demonstrating that the validity of Haff's law [3] is restricted to times less than one mean free time t_0 . This law states that the decay of the total energy from an initially homogeneous equilibrium state obeys $E(t) = E(0)/(1 + t/t_e)^2$, where $t_e = 2dt_0/(1 - \alpha^2)$ [3] in a d -dimensional IHS fluid, and α is the coefficient of normal restitution.
- (ii) An exact analytic relation for the time dependence of certain velocity moments, on the basis of which they question the validity of the kinetic theory of granular gases.

In the present comment I show that their analytic relation does not hold, and that their DSMC results refer to unphysical densities of twice close packing, and consequently have no physical relevance.

Concerning point (i): many authors have shown by means of molecular dynamics [4, 5], and of DSMC [6] that Haff's law is in quantitative agreement with simulation results for times up to the clustering transition (density instability), for all gas and fluid densities, and for small and moderate inelasticities with restitution coefficient $\alpha > 0.7$.

However, the authors of Ref. [2] arrive at quite different conclusions. From their input data for solving the two-dimensional Boltzmann equation by DSMC, I extract the following values, relevant for our discussion: number of particles, $N = 2 \times 10^4$; system size $L = 100\sigma$ (σ is a disk diameter) and mean free path $l_0 \simeq 9\sigma$. This yields the unphysical value for the coverage or area fraction $\phi = \frac{1}{4}\pi N\sigma^2/L^2 \simeq 1.57$, which is about twice (!) the closed packed density of hard disks. According to the Enskog theory the value $l_0 \simeq 9\sigma$ corresponds to a coverage of $\phi \simeq 0.03$, which is not consistent with the authors' data. Moreover Figure 1 and the text above Eq. (3) of Ref. [2] inform the reader that the mean free time is $t_0 \simeq 350dt$, where dt is the integration time step in DSMC, and that Haff's law breaks down at time $300dt \simeq (3/7)t_o$.

We conclude that the orders of magnitude of the simulation results in Ref. [2] have little relevance for DSMC and molecular dynamic simulations at realistic values of the system parameter.

Concerning point (ii): their criticism on granular kinetic theory for three-dimensional IHS fluid is based on the following incorrect relation for the velocity moments, $\partial_t \mu_k(t) = -2\pi n \sigma^2 (1 - \alpha^k) \mu_{k+1}(t)$, claimed to be valid in the translationally invariant homogeneous cooling state (HCS). The moments are defined as,

$$\mu_k(t) \equiv \langle |g_{12}|^k \rangle = \frac{1}{N(N-1)} \int dx_1 dx_2 |g_{12}|^k f_0^{(2)}(x_1 x_2 t). \quad (1)$$

Here $f_0^{(s)}(x_1 \dots x_s, t)$ are s -particle distribution functions which represent the probability density for finding an arbitrary set of s particles in the volume element $dx_1 \dots dx_s$, where $x_1 = \{\vec{r}_i, \vec{v}_i\}$ ($i = 1, 2, \dots, s$) in the HCS. These pdf's only depend on relative distances. Furthermore $g_{ij} \equiv \vec{v}_{ij} \cdot \hat{r}_{ij}$ with $\vec{v}_{ij} = \vec{v}_i - \vec{v}_j$, and $\hat{r}_{ij}$ is a unit vector along the lines of centres at arbitrary nonoverlapping positions inside the volume $V = L^3$ of the system. The moments $\mu_k(t)$, are intensive quantities, which approach for large V the values $(1/4\pi n^2) \times \int d\vec{v}_1 d\vec{v}_2 \int d\vec{r}_{12} |g_{12}|^k f_0(\vec{v}_1) f_0(\vec{v}_2)$, which are independent of the system size. This holds for arbitrary long range spatial correlations in $[f_0^{(2)}(12) - f_0(1)f_0(2)]$, which decay faster than

$1/r^a$ with $a > 0$. The above argument is made to establish that $\mu_k(t)$ is independent of the system size.

To calculate the rate of change $\partial_t \langle A(12) \rangle$ of an average pair variable $\langle A(12) \rangle$, as in Eq. (1), I use the second equation of the BBGKY-hierarchy for the reduced distribution functions in terms of binary collision operators $\bar{T}_{ij}$ as derived in Ref. [7, 8, 9], multiply it with $\langle A(12) \rangle / N^2$, integrate over x_1 and x_2 , and express it in the adjoint binary collision operator T_{ij} , which reads

$$T_{12} = \sigma^2 \int_{\vec{v}_{12} \cdot \hat{\sigma}_{12} < 0} d\hat{\sigma}_{12} |\vec{v}_{12} \cdot \hat{\sigma}_{12}| \delta(\vec{r}_{12} - \sigma \hat{\sigma}_{12}) (b_{\sigma_{12}} - 1). \quad (2)$$

Here $\hat{\sigma}_{12} = \hat{r}_{12}$ for two spheres in contact, and $b_{\sigma_{12}} A(\vec{r}_1 \vec{v}_1 \vec{r}_2 \vec{v}_2) = A(\vec{r}_1 \vec{v}_1^* \vec{r}_2 \vec{v}_2^*)$ with

$$\vec{v}_i^* = \vec{v}_i - \frac{1}{2}(1 + \alpha)(\vec{v}_{ij} \cdot \hat{\sigma}_{ij}) \hat{\sigma}_{ij}. \quad (3)$$

In fact, the explicit form of the operator T_{12} is not needed in the subsequent arguments, but only that the operator is non-vanishing when particles 1 and 2 are close together.

The result for the rate of change $\partial_t \langle A(12) \rangle$ is then:

$$\begin{aligned} \partial_t \langle A_{12} \rangle &= \frac{1}{N^2} \int dx_1 dx_2 f_0^{(2)}(x_1 x_2) \left(\vec{v}_{12} \cdot \frac{\partial}{\partial \vec{r}_{12}} + T_{12} \right) A_{12} + \\ &+ \frac{2}{N^2} \int dx_1 dx_2 dx_3 f_0^{(3)}(x_1 x_2 x_3) T_{13} A_{12} \\ &\equiv \mathcal{K}_1 + \mathcal{K}_{12} + \mathcal{K}_{123}. \end{aligned} \quad (4)$$

Here $\mathcal{K}_1$ and $\mathcal{K}_{12}$ represent the terms containing respectively $\vec{v}_{12} \cdot \partial / \partial \vec{r}_{12}$ and T_{12} on the first line of (4), and $\mathcal{K}_{123}$ the term on the second line. Then, taking $A(12) = |g_{12}|$, $\mathcal{K}_1$ vanishes, as shown in App.A of Ref. [2]. The $\{x_1, x_2\}$ -integration in $\mathcal{K}_{12}$ yields a factor of $\mathcal{O}(V)$, as $\delta(\vec{r}_{12} - \sigma \hat{\sigma}_{12})$ in (2) is constraint to the value $|\vec{r}_{12}| = \sigma$. Consequently, the contribution $\mathcal{K}_{12}$, introduced in (A7) of Ref. [2], is of $\mathcal{O}(1/V)$, and vanishes for large systems, and so do the RHS of Eqs. (A5) and (31) in Ref. [2].

This does not mean that $\partial_t \mu_k(t)$ in Eq. (31) of Ref. [2] should be vanishing. Inspection of $\mathcal{K}_{123}$ in (4) shows first that this term is $\mathcal{O}(1)$ for large V , because the $\{\vec{r}_1, \vec{r}_2\}$ -integrations yield a factor V^2 , because T_{13} poses the constraint $|\vec{r}_{13}| = \sigma$. For an explicit calculation I insert (2) in (4), and take the thermodynamic limit. This shows that the contribution of $[f_0^{(3)}(123) - f_0(2)f_0^{(2)}(13)]$ to $\mathcal{K}_{123}$ is of $\mathcal{O}(1/V)$. With the help of the collision dynamics (1) and (2) I arrive at the explicit result:

$$\begin{aligned} \mathcal{K}_{123} &= \frac{2\sigma^2}{4\pi n^2} \int d\vec{v}_1 d\vec{v}_2 d\vec{v}_3 \int'_{\vec{v}_{13} \cdot \hat{r}_{13} < 0} d\hat{r}_{12} d\hat{r}_{13} f_0(v_2) f_0^{(2)}(\sigma \hat{\sigma}_{13}, \vec{v}_1 \vec{v}_3) \\ &\times |\vec{v}_{13} \cdot \vec{r}_{13}| \left\{ |\vec{v}_{12} \cdot \hat{r}_{12} - \frac{1+\alpha}{2}(\vec{v}_{13} \cdot \hat{r}_{13}) \hat{r}_{13} \cdot \hat{r}_{12}|^k - |\vec{v}_{12} \cdot \hat{r}_{12}|^k \right\}. \end{aligned} \quad (5)$$

The integrations over $\{\vec{v}_1 \vec{v}_3 \hat{r}_{13}\}$ are restricted to the precollision hemisphere, $\vec{v}_{13} \cdot \hat{r}_{13} < 0$. Inspection shows that $\mathcal{K}_{123}$ is $\mathcal{O}(1)$ and nonvanishing, contrary to what is argued in Appendix A of Ref. [2].

In summary, the relation $\partial_t \mu_k(t) = -\Gamma_k \mu_{k+1}(t)$, for the velocity moments claimed to be exact, does not hold as it is based on an incorrect starting point, Eq. (31) of Ref. [2]. In deriving their Eq.(31) the authors retain a term $\mathcal{K}_{12}$ which is of $\mathcal{O}(1/V)$. This term should be neglected in comparison to $\partial_t \mu_k(t)$, which is $\mathcal{O}(1)$ for large V . Therefore, the main simulation and analytic results of this paper are incorrect.

References

- [1] H.H. Ernst and T.P.C. van Noije, *Kinetic Theory of Granular Gases*, Proceedings of the Bad Honnef Workshop on "Granular Gases", S. Luding and T. Pöschel (eds) (Springer Verlag, 1999).
- [2] J.M. Salazar and L. Brenig, Phys. Rev. E **59**, 2093 (1999); J. Plasma Phys. **59**, 639 (1998).
- [3] P.K. Haff, J. Fluid Mech. **134**, 401 (1983).
- [4] J.A.G. Orza, R. Brito, T.P.C. van Noije and M.H. Ernst, Int. J. Mod. Phys. C **8**, 953 (1997); T.P.C. van Noije, M.H. Ernst, R. Brito and J.A.G. Orza, Phys. Rev. Lett. **79**, 411 (1997); R. Brito and M.H. Ernst, Europhys. Lett. **43**, 497 (1998).
- [5] S. McNamara and W.R. Young, Phys. Rev. E **53**, 5089 (1996); P. Deltour and J.-L. Barrat, J. Phys. I France **7**, 137 (1997); S.E. Esipov and T. Pöschel, J. Stat. Phys. **86**, 1385 (1997); M. Müller and H.J. Herrmann, in *Physics of Dry Granular Media*, eds. H.J. Herrmann, J.-P. Hovi and S. Luding (Kluwer Academic Publishers, Dordrecht, 1998).
- [6] M. Müller and H.J. Herrmann; J.J. Brey, M.J. Ruiz-Montero and D. Cubero, Phys. Rev. E **54**, 3664 (1996); J.M. Montanero and A. Santos, private communication, March 1998.
- [7] M.H. Ernst, J.R. Dorfman, W.R. Hoegy and J.M.J. van Leeuwen, Physica **45**, 127 (1969).
- [8] J.J. Brey, J.W. Dufty and A. Santos, J. Stat. Phys. **87**, 1051 (1997).
- [9] T.P.C. van Noije, M.H. Ernst and R. Brito, Physica A **251**, 266 (1998).